

SA72B-0544 1330h POSTER

PML boundary conditions implementation in 1D and 2D wave propagation code for an inhomogeneous, magnetized ionospheric plasma.

N Gondarenko¹ (301-405-1621; natalia@glue.umd.edu)

P N Guzdar¹ (301-405-1590; guzdar@glue.umd.edu)

S L Ossakow² (202-767-2723; ossakow@ccs.nrl.navy.mil)

¹IREAP, University of Maryland, College Park, MD 20742

²Plasma Physics Division, Naval Research Laboratory, Washington, DC 20375

We have developed 1D and 2D full-wave propagation codes with arbitrary direction of propagation to the magnetic field and the plasma density relevant to ionospheric interaction studies. A wave launched from the bottom of computational box should be allowed to propagate out of the simulation box (top boundary for the 1D case and top and side boundaries for the 2D case) without providing spurious reflection from these computational boundaries. We have implemented the use of a Perfectly-Matched-Layer (PML) technique using matrix-marching algorithms to accomplish these goals. Results for various 1D and 2D cases with the existence of mode conversion layers for specified density profiles will be presented. These codes can be used to accurately compute the detailed mode structure of radio wave in complex magnetized plasma environments.

SA11A MCC: 134 Monday 0830h

Nicolet Lecture (joint with SH, SM)

Presiding: D N Baker, University of Colorado, Boulder

SA11A-01 0830h INVITED

Aeronomy: From Exploration to Data Assimilation

Robert W Schunk (435-797-2978; schunk@cc.usu.edu)

Utah State University, Center for Atmospheric and Space Sciences 4405 Old Main Hill, Logan, UT 84322-4405, United States

During the last century, there has been continual progress with regard to elucidating the important aeronomy processes that operate in the Earth's upper atmosphere. However, the progress was not uniform and there were certain periods of time when significant advances in knowledge were achieved due to an important discovery, a novel instrumental technique, or a new international program. Shortly after Marconi successfully transmitted radio signals across the Atlantic Ocean, the existence of the ionosphere was clearly established and its source, peak height, and layer characteristics were determined. In the mid-1940s, the rocket technology that became available was used by scientists to study the ionosphere and upper atmosphere. The great potential of this new technology, coupled with a major advance in ground-based instrumentation, led scientists to realize that a significant increase in knowledge of the terrestrial environment was possible. This resulted in the International Geophysical Year (1957-58) and the launch of the first satellite (Sputnik I). Another significant advance in aeronomy began in the mid-1980s, with the advent of supercomputers, the development of global numerical models, the existence of numerous ground-based instrument networks, and the proliferation of satellites. At the beginning of the new millennium, aeronomy is at the threshold of another rapid advance in knowledge due to the use of data assimilation techniques. Although data assimilation has been widely used by both meteorologists and oceanographers for several decades, only recently has there been a sufficient quantity of data for this approach to be useful in aeronomy. During the coming decade, tens of millions of measurements of the ionosphere-thermosphere system could become available on a daily basis from a myriad of in situ and remote sensing instruments. These data sets can be assimilated into a time-dependent, physics-based, numerical model of the ionosphere-thermosphere system via Kalman filters or other assimilation techniques. As a result, it will be possible to provide time-dependent 3-dimensional reconstructions of the ionospheric and thermospheric densities on an hourly basis day after day. These reconstructions will not only greatly advance aeronomy, but will also provide important societal benefits.

SA11B MCC: 134 Monday 1000h

Tracing the Sun-Earth Connection Into the Upper Atmosphere: Study of the April 2002 Events I (joint with SH, SM)

Presiding: L J Paxton, Applied Physics Laboratory; J U Kozyra, University of Michigan

SA11B-01 1000h INVITED

Tracing the Sun-Earth Connection: The April 14-24, 2002 Events

Richard Fisher (rfisher@hq.nasa.gov)

NASA Headquarters, Code S, Washington, DC 20546, United States

The interaction of the Earth with the Sun and the local space environment is complex and presents many challenges if that interaction is to be fully understood. The April 14-24, 2002 events may well serve as the catalyst for a new way of approaching the science that we do. A flotilla of spacecraft that can study the Sun, the interplanetary medium, the magnetosphere, and the Earth's atmosphere are now providing data. For the first time we can trace, in some detail, the chain of causality from the Sun to the Earth. We also have available to us a range of first principle models that allow us to validate our understanding of the physics of these processes by completely specifying their inputs and comparing their outputs to actual observations. Data provide the ultimate reality check on the models. This reality check will allow us to evaluate how faithfully models handle the transition from one region of applicability to another as we trace this storm from the sun to the Earth. For convenience we can consider this system of systems to consist of four chains that span physics disciplines: solar particles, magnetospheric drivers, solar radiance, and energy balance. In the invited talks that follow each element in this chain will be summarized. In this event we are able to trace the production of a disturbance on the Sun as active regions, flares and coronal mass ejections, through the interplanetary medium until it impinges upon the magnetosphere and interacts with the Earth's upper atmosphere. The April storm is unique, not only for the behavior of the storm, but for the addition of the information provided by TIMED as the newest element in the constellation of satellites providing data. We hope that this interactive and cooperative study will be a paradigm for future activities

URL: <http://storms.jhuapl.edu>

SA11B-02 1020h INVITED

Solar Radiance Chain in the April 2002 Series of Flares

Robert P Lin¹ (510-642-1149; rlin@ssl.berkeley.edu)

Stanley Solomon² (stans@ucar.edu)

¹Physics Dept Space Sciences Lab, University of California, Berkeley, University of California Space Sciences Laboratory, Berkeley, CA 94720-7450, United States

²HAO, NCAR, 3450 Mitchell Ln, Boulder, CO 80301, United States

The April 2002 series of solar storms included a number of GOES M- and X-class flares, in particular, a X1.5 flare on April 21 that was observed by many spacecraft - SoHO, TRACE, Wind, and the recently launched RHESSI (Ramaty High Energy Solar Spectroscopic Imager), which is designed to study flare hard X-ray/gamma-ray emissions. At the same time the effects of these emissions on the Earth were observed by a fleet of spacecraft, including TIMED, and by ground-based instrumentation. The effects of the increased energetic photon flux on the Earth's atmosphere during solar flares is usually small in comparison to geomagnetic disturbances, but large X-class flares (such as the one on April 21, 2002) that are several hours in duration can have significant consequences. When the 1 to 10 nm region of the solar spectrum is enhanced during a flare, the energy is deposited largely in the E-region of the ionosphere, 100 to 120 km altitude, where increases in ion density, photoelectron production, airglow emission, and odd-nitrogen production can result. Higher energy photons (such as observed by RHESSI) penetrate to lower altitudes, where they have less effect on the atmosphere but can still create additional ionization in the D-region of the ionosphere, 80 to 100 km altitude, that is disruptive of certain types of radio communication. Here we present the observations of solar photon emissions and their effects on the Earth's ionosphere and atmosphere.

URL: <http://storms.jhuapl.edu>

SA11B-03 1040h INVITED

Tracing the Sun-Earth Connection: The Solar Particle Chain in April 2002

D. N. Baker¹ (303-492-4509; daniel.baker@lasp.colorado.edu); R. A. Mewaldt² (dick@srll.caltech.edu); S. G. Kanekal³ (kanekal@surya.umd.edu); M. G. Henderson⁴ (mghenderson@lanl.gov); D. S. Evans⁵ (David.S.Evans@noaa.gov); S. M. Bailey⁶ (scott.bailey@gi.alaska.edu); M. J. Reiner⁷ (reiner@urap9.gsfc.nasa.gov); B. R. Dennis⁷ (Brian.R.Dennis.1@gsfc.nasa.gov)

¹Laboratory for Atmospheric and Space Physics/University of Colorado, 1234 Innovation Drive, Boulder, CO 80309, United States

²California Institute of Technology, MS 103-33, Pasadena, CA 91125, United States

³Dept. of Physics, Univ. of Maryland, College Park, MD 20742, United States

⁴Los Alamos National Laboratory, PO Box 1663, Los Alamos, NM 87545, United States

⁵NOAA, 325 S. Broadway, Boulder, CO 80307, United States

⁶Geophysical Institute, Univ. of Alaska, Fairbanks, AK 99775, United States

⁷Goddard Space Flight Center, Code 682, Greenbelt, MD 20771, United States

Soon after the RHESSI spacecraft observed an X-class solar flare early on 21 April 2002 a coronal mass ejection (CME) moving at ~2500 km/sec was observed by SOHO to emerge from near the west limb of the Sun. Almost simultaneously, Type-II and Type-III radio emission was observed by the Wind spacecraft, indicating the onset of particle acceleration near the Sun. The active region on the Sun responsible for this event was magnetically well-connected to the Earth, leading to a prompt enhancement at 1 A.U. of solar particles (>10 MeV) as seen by the ACE and SAMPEX spacecraft. Strong shock-related particle acceleration continued to occur as the CME propagated outward from the sun. The solar particle enhancement lasted from 21 April to 26 April (as seen by the NOAA/POES spacecraft) and included enhancement not only of solar protons and heavier nuclei extending to hundreds of MeV/nucleon but also solar electrons. The energetic particles produced near the sun, at this and preceding interplanetary shock waves, and deep within the Earth's radiation belts were all observed using TIMED and SNOE to produce substantial effects on the chemistry of the Earth's middle and upper atmosphere. We examine the impacts of this set of events on atmospheric heating and cooling rates and on ozone densities in the mesosphere. Using the remarkable constellation of available spacecraft, we are able to follow the particle chain from the Sun's surface all the way to the deep layers of Earth's atmosphere with unprecedented completeness.

SA11B-04 1100h INVITED

Overview of One Aspect of the Sun-Earth Connection during the April 2002 Events: the "Magnetospheric Driver" Chain

Thomas H Zurbuchen¹ ((734) 647-6835; thomasz@umich.edu); Janet Kozyra¹ (jukozyra@umich.edu); Gareth Lawrence² (gri@kreutz.nascom.nasa.gov); James L. Burch³ (jburch@swrl.org); Michael G. Henderson⁴ (mghenderson@lanl.gov); William J. Burke⁵ (william.burke@hanscom.af.mil); Larisa Goncharenko⁶ (lpg@haystack.mit.edu); James M. Russell⁷ (james.russell@hamptonu.edu); Raymond G. Roble⁸ (roble@ncar.ucar.edu)

¹University of Michigan, 2455 Hayward St, Ann Arbor, MI 48109-2143, United States

²NASA Goddard SFC, MC 682 3 Bldg 26/Rm G1, Greenbelt, MD 20771, United States

³Southwest Research Inst, 6220 Culebra Rd POD 28510, San Antonio, TX 78228, United States

⁴Los Alamos Natl Lab, MS D436, Los Alamos, NM 87545, United States

⁵Air Force Laboratory, 12 Vanderbilt Rd, Acton, MA 01720

⁶MIT, Haystack Observatory Route 40, Westford, MA 01886, United States

⁷Hampton University, Department of Physics, Hampton, VA 23668, United States

⁸NCAR, POB 3000, Boulder, CO 80307, United States