

suboxic sediments is well documented, DOC pore water gradients have most often been interpreted uniquely as indicative of a flux of DOC out of the sediments to the overlying bottom waters [2-5]. This interpretation stems from our inability to resolve a subsurface DOC sink from the vertical distribution of sedimentary particulate organic carbon (POC). This DOC sink, if it exists, would considerably alter our views of the mechanisms that regulate DOC fluxes across the sediment-water interface as well as their quantification. Sorption onto authigenic metal oxides may also lead to a molecular and isotopic fractionation of DOC. Furthermore, if DOC sorption to metal oxides is a reversible process, it may buffer pore water DOC concentrations and release to the overlying waters. Oxidic surface sediments recovered from the St. Lawrence Estuary and the Saguenay Fjord were incubated under anaerobic conditions and extracted with a mild reducing agent to determine the amount and composition (molecular and isotopic) of the DOC associated with the authigenic iron and manganese oxides. Preliminary results from the study will be presented. References: [1] Deflandre et al. (2002) *Geochim. Cosmochim. Acta*, 66; 14, 2547-2558. [2] Alperin et al., (1999) *Geochim. Cosmochim. Acta*, 63; 3-4, 427-448. [3] Burdige et al., (1999) *Geochim. Cosmochim. Acta*, 63; 10, 507-515. [4] Holcombe et al., (2001) *Limnol. Oceanogr.*, 46; 2, 298-308. [5] Papatimitriou et al. (2002) *Mar. Chem.*, 79, 37-47.

OS43A-07 1330h POSTER

Contributions of Wind Forcing and Surface Heating to Interannual Sea Surface Height Variability in the Atlantic Ocean.

Cecile Cabanes¹ (33-2-98-22-42-95; cecile.cabanes@ifremer.fr)

Thierry Huck² (33-2-98-01-65-10; thuck@univ-brest.fr)

Alain Colin de Verdiere² (33-2-98-01-62-24; acolindv@univ-brest.fr)

¹Laboratoire de Physique des Océans, Ifremer, centre de Brest B.P. 70, Plouzané 29280, France

²Laboratoire de Physique de Océans, UBO, UFR Sciences 6, avenue Le Gorgeu B.P. 809, Brest 29285, France

Interannual sea surface height variations in the Atlantic Ocean are examined from 10 years of high-precision altimeter data in light of simple mechanisms that describe the ocean response to atmospheric forcing: (1) local steric changes due to surface buoyancy forcing and a local response to wind stress via Ekman pumping, and (2) baroclinic and barotropic oceanic adjustment via propagating Rossby waves and quasi-steady Sverdrup balance respectively. It is shown that most of the interannual sea level variability is related to local response to heat fluxes changes (more than 50% in the eastern North Atlantic). Except in a few places, a local response to wind stress forcing is less successful in explaining sea surface height observations. In this case, it is necessary to consider large scale oceanic adjustments: first baroclinic mode forced by wind stress explain more than 30% of interannual sea level variations in the eastern North Atlantic or in the South Atlantic (between 30 and 40 ° S) and in the range of 70 % in the latitude band 17-20 ° N. A quasi-steady sverdrup response is also observed between 40 and 45 ° N.

OS43A-08 1330h POSTER

Multi-Platform Seabed Classification of Georgia Strait, British Columbia (Canada)

Liliane Carle¹ (lcarle@uvic.ca)

Stephen F. Bloomer¹ (sbloomer@uvic.ca)

Ross N Chapman² (chapman@uvic.ca)

¹C-MARS, University of Victoria, P.O. Box 3055, Victoria, BC V8W 3P6, Canada

²School of Earth and Ocean Sciences, University of Victoria, P.O. Box 3055, Victoria, BC V8W 3P6, Canada

This paper presents the results of a multi-platform seabed classification in Georgia Strait. This study is part of an NSERC CRD project to establish a methodology to efficiently develop a comprehensive understanding of the seabed, based on the integration of the classification of acoustic data from multiple sonar platforms with suitable auxiliary groundtruth data. Two different data sets are classified and integrated in this paper. Classification of the backscatter of Simrad EM1002 multibeam data generated a high spatial resolution map related to the seafloor geology. Classification of high-resolution Huntce seismic lines in the same area produced a map of the subbottom seabed geology that can be compared to the multibeam seafloor classification map. The main challenge addressed in

this paper is that EM1002 often operates in 3 discrete modes to optimize the quality of the bathymetric data. Generally these modes are depth related, with a longer acoustic pulse used in areas where the seafloor is deeper. For classification purposes, this means that each mode has to be treated separately, because the pulse length strongly influences the backscatter statistics. From here, a general multibeam classification was developed from the 3 separate classification results and integrated with the seismic classification results.

OS43A-09 1330h POSTER

Testing the Reliability and Sensitivity of Foraminiferal Transfer Functions Based on the Modern Analog Technique (MAT)

Don Lac¹ (1-978-542-6282; donl1214@hotmail.com)

James L Cullen¹ (1-978-542-6283; james.cullen@salemstate.edu)

April Martin²

¹Department of Geological Sciences, Salem State College, Salem, MA 01970, United States

²Department of Geological Sciences, Brown University, Providence, RI 02912, United States

Quantitative estimates of past sea-surface temperatures (SST's) based on surface sediment calibration data sets of planktic foraminifers and modern SST's have been widely used in the interpretation and modeling of past climates. One widely used approach, The Modern Analog Techniques (MAT) relies on comparing a downcore sample to the Brown University modern Global Data Base of 1265 seabed samples and choosing either the 10 or 5 most similar modern samples using the squared-chord distance similarity metric. The SST's above the best modern analogs are then averaged to produce the downcore SST estimate. We have chosen a set of 8 modern sea-bed samples from the Global Data Base with a wide range of foraminiferal compositions; 3 from the Pacific, 3 from the Atlantic, and 2 from the Indian Ocean and have generated duplicate foraminiferal census counts from sets of 5-6 random splits from each of our 8 samples so that we can: 1. compare the degree of similarity between duplicate samples so that we can evaluate the differences in dissimilarity values that can be attributed to counting error and begin to better understand the sensitivity of the chosen dissimilarity measures to ecologically produced differences in foraminiferal composition, 2. evaluate differences in how the duplicate samples choose analogs from the Global Data Base, and 3. test the sensitivity of the MAT's ability to accurately and precisely predict SST's using analogs from the Global Data Base for each set of duplicate samples. Comparison of the dissimilarity coefficients within each set of duplicate samples produces maximum dissimilarity values that range from 0.03 to 0.14. Both mean and maximum dissimilarities are greatest in sample sets from the low latitudes. The 5 best analogs chosen from the Global Data Base for samples within each set of duplicates generate average dissimilarities that range from 0.01 to 0.04. However, between a total of 8 and 12 different modern analogs were needed to find the 5 best analogs for all the samples within a particular set of duplicates, with no regional pattern in this number observed. The warm and cold SST estimates generated using the SST's above each of the 5 chosen analogs exhibit a wide range of variation, particularly for the three sample sets from the high latitudes. The three subpolar sample sets exhibit a 3.4, 1.1, and 1.0 degree C range in their cold SST estimates. There is no clear relationship between differences in SST estimates and differences in the average dissimilarities within duplicate sample sets. Using 10 instead of 5 modern analogs to estimate SST's produces somewhat better results for 4 of the 8 sample sets and similar results for the remaining 4. Our results suggest that foraminiferal samples with dissimilarity values of up to 0.15 are not detectably different from duplicate foraminiferal census counts and should be considered excellent modern faunal analogs for any fossil sample. In addition high latitude samples seem to be produce somewhat less reliable SST estimates than low latitude samples. Finally, our results suggest that, when estimating past SST's choosing to average the SST's above the 10 best analogs produces more accurate (and precise) results, particularly in situations where the Global Data Base contains adequate modern analogs.

OS44A CC: 520 A Thursday 1530h

Rachel Carlson Lecture (joint with U)

Presiding: J A Carton, University of Maryland

OS44A-01 1535h

Beyond Correlations: the Search for Mechanisms Underlying Coupled Climate/Ecosystem Variability in the Oceans

Ann E Gargett (757-683-6009; gargett@ccpo.edu.edu)

Ann Gargett, Center for Coastal Physical Oceanography Dept of Ocean, Earth and Atmospheric Sciences Old Dominion University, Norfolk, VA 23508, United States

Correlations between variables describing some part of the earth's climate system and others associated with some part of oceanic ecosystems are widely used to suggest the existence of causal linkages. I will consider the dangers inherent in using such correlations for predictive purposes, and argue that real predictive skill demands identification of the underlying connective mechanisms. As an example, I will describe a possible mechanistic explanation for observed correlations between the winter state of the atmosphere over the North Pacific Ocean and the success of various North Pacific salmon fisheries. This same context will be used to demonstrate that the crucial final step, that of testing candidate mechanisms, is presently severely limited by lack of long-term data sets that include biological variables other than fish catches. Identification and understanding of the mechanisms connecting variability in climate and large marine ocean ecosystems is both economically essential, and increasingly urgent as the earth's climate system moves into states outside the experience of observational earth sciences in the modern age.

OS51A CC: 524 C Friday 0830h

General Ocean Sciences II

Presiding: H Sun, Rider University; J E Friddell, ERDC-Cold Regions Research and Engineering Laboratory

OS51A-01 0830h

Decadal Changes in the North Atlantic Subpolar Circulation

Sirpa Hakkinen¹ (301-614-5712; sirpa.hakkinen@nasa.gov)

Peter B Rhines² (206-543-0593; rhines@ocean.washington.edu)

¹NASA Goddard Space Flight Center, Code 971, Greenbelt, MD 20771, United States

²University of Washington, Box 357940, Seattle, WA 98195, United States

Observations of sea surface height reveal that dramatic changes have occurred over the past decade in the mid-to-high-latitude North Atlantic Ocean. Altimeter data from TOPEX/Poseidon mission show that subpolar sea surface height increased during the 1990s and the geostrophic velocity derived from altimeter data shows a decline in the subpolar gyre circulation. Combining the data from earlier satellites, Seasat (1978) and Geosat (1985-1988), we find that at the late 1990s the subpolar circulation may actually be considerably weaker than in the (late 1970s and) 1980s. Direct current-meter observations in the boundary current of the Labrador Sea support the circulation trend of the 1990s and, together with hydrographic data, show that the mid-to-late 1990s decline extends deep in the water column. Based on analysis of the local surface forcing we find that buoyancy forcing over the northern North Atlantic has a dynamic effect consistent with the altimeter data and hydrographic observations: a weak thermohaline forcing and the subsequent decay of the domed structure of the subpolar isopycnals would support the observed decline of the gyre circulation. Only continued remote and in situ observations will show whether this 1990s change is temporary or part of much longer term variability.

OS51A-02 0845h

Subantarctic Mode Water Formation South of Australia

Paul Barker^{1,2} (paul.barker@acadiau.ca)

John Bye² (j.bye@earthsci.unimelb.edu.au)

¹Acadia university, Dept of maths and Stats Acadia university, Wolfville, NS b4p2r6, Canada

²University of Melbourne, Earth Sciences University of Melbourne, Parkville, Vic 3010, Australia

Subantarctic Mode Water (SAMW) occurs north of the Antarctic Circumpolar Current (ACC) in the Southern Ocean, where it is characterised by its homogenous properties. A new method is presented for classifying and identifying SAMW. From this it is possible that SAMW is formed south-west of Tasmania by a mixture of higher salinity and temperature water originating in the Tasman Sea, and lower salinity and temperature water originating in the Southern Ocean. The SAMW spreads from the source region into the South Australian Basin as a plume. A modulation in the two source water masses could account for the change in water properties in the upper 1 km of the South Australian Basin, where a warming of 1C has been observed since 1970.

OS51A-03 0900h

Using the Stable Isotopes of Carbon as a Tracer for Deep Ocean Circulation Across the Paleocene-Eocene Boundary

Flavia Nunes¹ (858-822-4925; fnunes@ucsd.edu)

Richard Norris¹ (858-822-1868; rnorris@ucsd.edu)

¹Scripps Institution of Oceanography, 9500 Gilman Drive MC 0208, La Jolla, CA 92092, United States

The Paleocene-Eocene (P/E) boundary is a time of abrupt fluctuations in the stable isotope record. In particular, the boundary is marked by a sharp decline to lighter values in carbon isotope ratios in various sites throughout the world. In order to assess how the carbon isotope distribution in the deep ocean varied before, during and after the event, a compilation of all existing carbon isotope data from benthic foraminifera across the P/E boundary, as well as high resolution data from 5 new sites has been assembled. The distribution of $\delta^{13}C$ in the modern oceans follows the pattern of deep ocean circulation, where water masses recently in contact with the atmosphere have a heavy isotopic ratio compared to low isotopic ratios in aged water masses. Assuming that the processes responsible for this distribution pattern also occurred at the time of the P/E boundary, then $\delta^{13}C$ can be used as a tracer for deep ocean circulation. Data from 12 deep-sea sites suggest that water mass aging gradients at this time appear to be much smaller than Late Paleocene or modern values. This may be a result of increasing global temperatures, which may be responsible for the weakening of deep water formation at higher latitudes. Decreased export production may also provide an explanation, since the lower availability of organic matter would decrease the source of light carbon being incorporated in aging water masses.

OS51A-04 0915h

Investigation of the Physics of Large Step-like Structures

Theodore D. Foster (1-858-822-2478; tfoster@seadog.ucsd.edu)

Scripps Institution of Oceanography, University of California, San Diego, La Jolla, CA 92093-0220, United States

Since 1969, on our many expeditions to the Southern Ocean we have observed large (from about 10 to over 100 meters thick) step-like structures in the vertical profiles of temperature and salinity at depths from about 100 to about 600 meters, but so far have only observed these phenomena in the central Weddell Sea. In order to study the physical cause of these large step-like structures, we have carried out several time series of measurements using CTD lowerings at regular time intervals while the ship drifted with the ice pack at various locations around Antarctica. From these time series of CTD profiles in the central Weddell Sea we found these large steps are carried up and down with internal tides, which would change pressures at interfaces sufficiently to cause instabilities. This effect might enhance the convection that may be the cause of these very large steps. The "density ratio" at the interfaces of these large steps always seems to be about 1.0, while for the smaller steps observed in the thermocline here it is about 1.4 and near the surface of the Arctic Ocean about 6 to 7. Somewhat similar large step-like structures have been observed recently by the Canadians in the Canada Basin of the Arctic Ocean at depths below about 2400 m. The cause of these steps may involve some of the same physical phenomena, but unless the renewal of the deep waters in the basin is rather rapid, the physical situation is very different than in the central Weddell Sea where the Ekman drift probably causes significant upwelling.

OS51A-05 0930h

Inter-Annual Variability of the St. Lawrence Productivity: a Modeling Study

Marina Chifflet¹ (1-418-775-0837; ChiffletM@dfo-mpo.gc.ca)

Vincent Le Fouest² (1-418-723-1981873; vincent_lefouest@uqar.qc.ca)

Michel Starr¹ (1-418-775-0611; StarrM@dfo-mpo.gc.ca)

François-Jean Saucier^{1,2} (1-418-723-1986#1328; SaucierF@dfo-mpo.gc.ca)

Bruno Zakardjian² (1-418-723-1986#1570; bruno.zakardjian@uqar.qc.ca)

¹Maurice-Lamontagne Institute, Fisheries and Oceans Canada, 850, route de la Mer, Mont-Joli, QC G5H 3Z4, Canada

²Institut des Sciences de la Mer (ISMER), Université du Québec à Rimouski, 310, allée des Ursulines, Rimouski, QC G5L 3A1, Canada

The productivity of the St. Lawrence is strongly dependent on circulation and fresh water input. How natural climate variability or climate changes may affect the dynamics of this ecosystem need to be evaluated in order to improve its predictability. Here, we present results of a coupled 3D physical-ecosystem model of the Estuary and Gulf of St. Lawrence that was developed to study the complex interactions between climate forcing and the productivity of biological resources at the base

of the food chain. The physical model is a high resolution 3D baroclinic circulation model (5 km in the horizontal and 5 m in the vertical) that is presently operational for the St. Lawrence basin (Saucier et al. 2003). It includes a level 2.5 turbulent kinetic energy scheme and an ice module. The ecosystem model includes eight compartments: nitrate and ammonium, two classes of phytoplankton (diatoms and flagellates), two classes of zooplankton (microzooplankton and mesozooplankton), and two sizes of detritus (Le Fouest et al. 2004). This ecosystem model is driven by the physical model using realistic tidal, atmospheric and hydrologic forcing for three years: 1997 as a typical year with high ice cover and high runoff, 1999 as a warm and productive year with low runoff and low ice cover and 1998, an intermediate year with low ice cover and high runoff. The simulated results of the seasonal and inter-annual variability in ocean climate and primary production are compared to field observations from the Atlantic Zonal Monitoring Program. Particular attention is directed towards the influence of climatic and hydrologic factors that exert a control on the nutrient distribution, seasonal plankton biomass, timing and magnitude of the spring bloom and primary production. Our results highlight the strong influence of winter hydrographic processes on the inter-annual variability in the St. Lawrence productivity. Le Fouest, V., Zakardjian, B., Saucier, F. J., Starr, M. 2004. Physical and climatic induced subregional variability of planktonic production in a high latitude marginal sea: the Gulf of St. Lawrence (Canada), in preparation. Saucier, F. J., Roy, F., Gilbert, D., Pellerin, P., Ritchie H. 2003. Modeling the formation and circulation processes of water masses and sea ice in the Gulf of St. Lawrence, Canada. J. Geophys. Res., 108(C8):1-20.

OS51A-06 0945h

The Connection Between Holocene Sea-Level Change in the Bering Sea-North Pacific and Beach-Ridge Genesis on Kamchatka's Eastern Coastline

Charles Henry Thibault (206-427-4137; zandar96@u.washington.edu)

University of Washington, Department of Earth Sciences 711e Condon Hall, Seattle, WA 98195, United States

The purpose of this study is to develop a Holocene sea level curve for the western North Pacific-Bering Sea region by utilizing information contained within profiles and stratigraphy of beach ridge sets on the eastern coast of Kamchatka, Russia. In order to successfully develop such a curve it is first necessary to find a site that is dominated by oceanic processes and to eliminate sources of noise by processes such as tectonics, isostasy, eolian transport, and fluvial and volcanic sediment fluxes. Each of these processes can influence the genesis of beach ridges, and each process contributes specific structural components to the overall morphology of a beach ridge set. Over the past seven years an international collaboration of scientists has investigated the eastern coast of Kamchatka, in part to study the relationship between numerous coastal processes and Kamchatka's enigmatic coastline. During this collaboration, beach ridge profile and stratigraphy data have been collected from seven study areas located along 1000 km of the eastern coast of Kamchatka. The study reported herein focuses on three sites: Khodutka, Ozernoi, and specifically on Uka which is a site apparently controlled by oceanic processes and which can therefore be utilized in the development of a Holocene sea level curve.